Observation of

Coherent Elastic Neutrino-Nucleus Scattering

Kate Scholberg, Duke University

Fermilab Wine&Cheese

September 1, 2017

OUTLINE

- Coherent elastic neutrino-nucleus scattering (CEvNS)
- Why measure it? Physics motivations (short and long term)
- How to measure CEvNS
- The COHERENT experiment at the SNS
- **First light** with CsI[Tl]
- Status and prospects for COHERENT

OUTLINE

- **Coherent elastic neutrino-nucleus scattering (CEvNS)**
- Why measure it? Physics motivations (short and long term)
- How to measure CEvNS
- The COHERENT experiment at the SNS
- **First light** with CsI[Tl]
- Status and prospects for COHERENT

Coherent elastic neutrino-nucleus scattering (CEvNS)

$$\nu + A \rightarrow \nu + A$$



A neutrino smacks a nucleus via exchange of a Z, and the nucleus recoils as a whole; **coherent** up to E_ν~ 50 MeV



Nucleon wavefunctions in the target nucleus are **in phase with each other** at low momentum transfer

$$\frac{d\sigma}{d\Omega} \sim A^2 |f(\mathbf{k}', \mathbf{k})|^2$$

Momentum transfer $\mathbf{Q} = \mathbf{k}' - \mathbf{k}$

For $QR << 1$,

[total xscn] ~ A^2 * [single constituent xscn ($|f|^2$)]

First proposed 43 years ago!

PHYSICAL REVIEW D VOLUME 9, NUMBER 5 1 MARCH 1974

Coherent effects of a weak neutral current

Daniel Z. Freedman†
National Accelerator Laboratory, Batavia, Illinois 60510
and Institute for Theoretical Physics, State University of New York, Stony Brook, New York 11790
(Received 15 October 1973; revised manuscript received 19 November 1973)

Our suggestion may be an act of hubris, because the inevitable constraints of interaction rate, resolution, and background pose grave experimental difficulties for elastic neutrino-nucleus scattering. We will discuss these problems at the end of this note, but first we wish to present the theoretical ideas relevant to the experiments.

Also: D. Z. Freedman et al., “The Weak Neutral Current and Its Effect in Stellar Collapse”, Ann. Rev. Nucl. Sci. 1977. 27:167-207

This is ***not*** coherent pion production,
a strong interaction process ***(inelastic)***





A. Higuera et. al, MINERvA collaboration,
PRL 2014 113 (26) 2477

\begin{aside}

Literature has CNS, CNNS, CENNS, ...

- I prefer including “E” for “elastic”... otherwise it gets frequently confused with coherent pion production at ~GeV neutrino energies
- I’m told “NN” means “nucleon-nucleon” to nuclear types
- CEνNS is a possibility but those internal Greek letters are annoying

➔**CEvNS**, pronounced “sevens”...

spread the meme!

\end{aside}

The cross section is cleanly predicted in the Standard Model

$$\frac{d\sigma}{dT} = \frac{G_F^2 M}{\pi} F^2(Q) \left[(G_V + G_A)^2 + (G_V - G_A)^2 \left(1 - \frac{T}{E_\nu}\right)^2 - (G_V^2 - G_A^2) \frac{MT}{E_\nu^2} \right]$$

E_ν: neutrino energy
T: nuclear recoil energy
M: nuclear mass
Q = √ (2 M T): momentum transfer

G_V, G_A: SM weak parameters

vector $G_V = g_V^p Z + g_V^n N,$ ← dominates

axial $G_A = g_A^p (Z_+ - Z_-) + g_A^n (N_+ + N_-)$ ← small for most nuclei, zero for spin-zero

$$g_V^p = 0.0298$$
$$g_V^n = -0.5117$$
$$g_A^p = 0.4955$$
$$g_A^n = -0.5121.$$

The cross section is cleanly predicted in the Standard Model

$$\frac{d\sigma}{dT} = \frac{G_F^2 M}{\pi} F^2(Q) \left[(G_V + G_A)^2 + (G_V - G_A)^2 \left(1 - \frac{T}{E_\nu}\right)^2 - (G_V^2 - G_A^2)\frac{MT}{E_\nu^2} \right]$$

E_ν: neutrino energy
T: nuclear recoil energy
M: nuclear mass
Q = √ (2 M T): momentum transfer

$F^2(Q)$: nuclear form factor, <~5% uncertainty on event rate



form factor suppresses cross section at large Q

For T<<E_ν, neglecting axial terms:

$$\frac{d\sigma}{dT} = \frac{G_F^2 M}{2\pi} \frac{Q_W^2}{4} F^2(Q) \left(2 - \frac{MT}{E_\nu^2}\right)$$

$$Q_W = N - (1 - 4\sin^2\theta_W)Z$$: weak nuclear charge

$\sin^2\theta_W = 0.231$,
so protons unimportant

$$\Rightarrow \boxed{\frac{d\sigma}{dT} \propto N^2}$$




The cross-section is *large*
^{133}Cs CEvNS
^{127}I CEvNS
ν_e ^{127}I CC
IBD
ν_e-e
(per target atom in CsI)
Cross-section (10^{-38} cm^2)
10
1
10^{-1}
10^{-2}
10^{-3}
5
10
15
20
25
30
35
40
45
50
55
Neutrino Energy (MeV)

Large cross section (by neutrino standards) but hard to observe due to **tiny nuclear recoil energies:**



➔ but **WIMP dark matter detectors** developed over the last ~decade are sensitive to ~ keV to 10's of keV recoils

The only experimental signature:

tiny energy deposited by nuclear recoils in the target material



OUTLINE

- Coherent elastic neutrino-nucleus scattering (CEvNS)
- **Why measure it? Physics motivations (short and long term)**
- How to measure CEvNS
- The COHERENT experiment at the SNS
- **First light** with CsI[Tl]
- Status and prospects for COHERENT

CEvNS: what's it good for?

1 So
2 Many
3 Things

!

(not a complete list!)

- Dark matter direct-detection background
- Well-calculable cross-section in SM:
  - $\sin^2\theta_{Weff}$ at low Q
  - **Probe of BSM physics**
    - Non-standard interactions of neutrinos
    - New NC mediators
    - Neutrino magnetic moment
- New tool for sterile neutrino oscillations
- Astrophysical signals (solar & SN)
- Supernova processes
- Nuclear physics:
  - Neutron form factors
  - g_A quenching
- Possible applications (reactor monitoring)









CEvNS from natural neutrinos creates ultimate background for direct DM search experiments

J. Billard, E. Figueroa-Feliciano, and L. Strigari, arXiv:1307.5458v2 (2013).



Understand nature of background (& detector response, DM interaction)

Clean SM prediction for the rate ➔ measure $\sin^2\theta_W$eff ; **deviation probes new physics**

$$\sigma \sim \frac{G_f^2 E^2}{4\pi}(N - (1 - 4\sin^2\theta_W)Z)^2$$



Example: hypothetical dark Z mediator (explanation for g-2 anomaly)

CEvNS sensitivity is @ low Q;
need sub-percent precision to compete w/ electron scattering & APV, but **new channel**

Non-Standard Interactions of Neutrinos:

new interaction **specific to ν's**

$$\mathcal{L}_{\nu H}^{NSI} = -\frac{G_F}{\sqrt{2}} \sum_{\substack{q=u,d \\ \alpha,\beta=e,\mu,\tau}} [\bar{\nu}_\alpha \gamma^\mu (1-\gamma^5)\nu_\beta] \times (\varepsilon_{\alpha\beta}^{qL}[\bar{q}\gamma_\mu(1-\gamma^5)q] + \varepsilon_{\alpha\beta}^{qR}[\bar{q}\gamma_\mu(1+\gamma^5)q])$$



J. Barranco et al., JHEP 0512 (2005), K. Scholberg, PRD73, 033005 (2006), 021

Can improve ~order of magnitude beyond CHARM limits with a first-generation experiment (for best sensitivity, want ***multiple targets***)

More studies: see https://sites.duke.edu/nueclipse/files/2017/04/Dent-James-NuEclipse-August-2017.pdf

Oscillations to sterile neutrinos w/CEvNS

(NC is flavor-blind): a potential new tool;

look for deficit and spectral distortion vs L,E

Examples:



Multi-πDAR sources at different baselines (20 & 40 m)



B. Dutta et al, arXiv:1511.02834

Anderson et al., PRD86 (2012) 013004, arXiv:1201.3805

Neutrino magnetic moment

Signature is **distortion at low recoil energy E**

$$\left(\frac{d\sigma}{dT}\right)_m = \frac{\pi\alpha^2\mu_\nu^2 Z^2}{m_e^2}\left(\frac{1-T/E_\nu}{T}+\frac{T}{4E_\nu^2}\right)$$



➔requires low energy threshold

See also Kosmas et al., arXiv:1505.03202

Nuclear physics with CEvNS

If systematics can be reduced to ~ few % level,
we can start to explore nuclear form factors

P. S. Amanik and G. C. McLaughlin, J. Phys. G 36:015105

K. Patton et al., PRC86 (2012) 024612

$$\frac{d\sigma}{dT} = \frac{G_F^2 M}{2\pi} \frac{Q_W^2}{4} F^2(Q) \left(2 - \frac{MT}{E_\nu^2}\right)$$

Form factor: encodes information about nuclear (primarily neutron) distributions

Fit recoil ***spectral shape*** to determine the F(Q^2) moments

(requires very good energy resolution,good systematics control)

Example:
tonne-scale
experiment
at πDAR source



Also note: tonne-scale low-threshold underground can look at **astrophysical neutrinos**

Solar neutrinos

J. Billard et al.,
Phys.Rev. D91 (2015) no.9, 095023



Rule out sterile oscillations using CEvNS (NC),
10 ton-year of Ge

R. Harnik et al., JCAP 1207 (2012) 026



Effect of new physics on CEvNS recoil spectrum

Supernova neutrinos in tonne-scale DM detectors



~ handful of events per tonne @ 10 kpc: sensitive to ***all flavor components*** *of the flux*

OUTLINE

- Coherent elastic neutrino-nucleus scattering (CEvNS)
- Why measure it? Physics motivations (short and long term)
- **How to measure CEvNS**
- The COHERENT experiment at the SNS
- **First light** with CsI[Tl]
- Status and prospects for COHERENT

How to detect CEvNS?



You need a neutrino source
and a detector

What do you want for your ν source?

- ✓ High flux
- ✓ Well understood spectrum
- ✓ Multiple flavors (physics sensitivity)
- ✓ Pulsed source if possible, for background rejection
- ✓ Ability to get close
- ✓ Practical things: access, control, ...

Both **cross-section and maximum recoil energy increase with neutrino energy**:





Want energy as large as possible while satisfying coherence condition: $Q \lesssim \frac{1}{R}$ (<~ 50 MeV for medium A)

Stopped-Pion (πDAR) Neutrinos





$$\pi^+ \longrightarrow \mu^+ + \nu_\mu$$

2-body decay: monochromatic 29.9 MeV ν_μ
PROMPT

$$\downarrow$$

$$\mu^+ \longrightarrow e^+ + \bar{\nu}_\mu + \nu_e$$

3-body decay: range of energies between 0 and $m_\mu/2$
DELAYED (2.2 μs)

Comparison of pion decay-at-rest ν sources

from duty cycle



$\propto \nu$ flux



Proton beam energy: 0.9-1.3 GeV
Total power: 0.9-1.4 MW
Pulse duration: 380 ns FWHM
Repetition rate: 60 Hz
Liquid mercury target

Time structure of the SNS source

60 Hz *pulsed* source



Background rejection factor ~few x 10^{-4}

The SNS has **large, extremely clean** DAR ν flux

0.08 neutrinos per flavor per proton on target



SNS flux (1.4 MW):
430 x 10^5 ν/cm²/s
@ 20 m

Now, ***detecting*** the tiny kick of the neutrino...

This is just like the tiny thump of a WIMP;
we benefit from the last few decades of low-energy nuclear recoil detectors



- low background (although for beam, requirements less stringent than for WIMPs)
- low energy threshold
- energy resolution
- fast timing
- nuclear recoil discrimination
- well-known (and large if possible) **quenching factor**
  (fraction of observable energy, keVr = QF* keVee)

Backgrounds

Usual suspects:

- cosmogenics
- ambient and intrinsic radioactivity
- detector-specific noise and dark rate

Neutrons are especially not your friends*

(although they sometimes give you a hand with calibration)



Steady-state backgrounds can be *measured* off-beam-pulse
... in-time backgrounds must be carefully characterized

*Thanks to Robert Cooper for the “mean neutron”

A “friendly fire” in-time background: **Neutrino Induced Neutrons (NINs)**

$$\nu_e + {}^{208}\mathrm{Pb} \rightarrow {}^{208}\mathrm{Bi}^* + e^- \quad \mathrm{CC}$$

$\searrow$ 1n, 2n emission

$$\nu_x + {}^{208}\mathrm{Pb} \rightarrow {}^{208}\mathrm{Pb}^* + \nu_x \quad \mathrm{NC}$$

$\searrow$ 1n, 2n, γ emission



- potentially non-negligible background from shielding
- requires careful shielding design
- large uncertainties (factor of few) in xscn calculation
- [Also: a signal in itself, e.g, HALO SN detector]



OUTLINE

- Coherent elastic neutrino-nucleus scattering (CEvNS)
- Why measure it? Physics motivations (short and long term)
- How to measure CEvNS
- **The COHERENT experiment at the SNS**
- **First light** with CsI[Tl]
- Status and prospects for COHERENT

The COHERENT collaboration

http://sites.duke.edu/coherent



COHERENT

SNS SPALLATION NEUTRON SOURCE

~80 members,
18 institutions
4 countries

arXiv:1509.08702

THE UNIVERSITY OF CHICAGO · Duke UNIVERSITY · UF UNIVERSITY of FLORIDA · KAIST · Laurentian University Université Laurentienne · Los Alamos NATIONAL LABORATORY EST. 1943 · BERKELEY LAB · MEPhI · NORTH CAROLINA CENTRAL UNIVERSITY FOUNDED 1910 · NC STATE UNIVERSITY · NM STATE UNIVERSITY · OAK RIDGE National Laboratory · Sandia National Laboratories · THE UNIVERSITY of TENNESSEE KNOXVILLE · TUNL · UNIVERSITY of WASHINGTON

U.S. DEPARTMENT OF ENERGY · Office of Science · NSF · KICP · NNSA National Nuclear Security Administration · CNEC

COHERENT CEvNS Detectors

Nuclear Target	Technology	Mass (kg)	Distance from source (m)	Recoil threshold (keVr)
CsI[Na]	Scintillating Crystal	14.6	19.3	6.5
Ge	HPGe PPC	10	22	5
LAr	Single-phase	22	29	20
NaI[Tl]	Scintillating crystal	185*/ 2000	28	13

Multiple detectors for N^2 dependence of the cross section





Siting for deployment in SNS basement

(measured neutron backgrounds low,
~ 8 mwe overburden)

View looking
down "Neutrino Alley"





Isotropic ν glow from Hg SNS target

Neutron Backgrounds

Several background measurement campaigns have shown that Neutrino Alley in the basement is neutron-quiet

SciBath



Sandia scatter cam



Expected recoil energy distribution



Prompt defined as first μs; note some contamination from ν_e and ν_μ-bar

OUTLINE

- Coherent elastic neutrino-nucleus scattering (CEvNS)
- Why measure it? Physics motivations (short and long term)
- How to measure CEvNS
- The COHERENT experiment at the SNS
- **First light with CsI[Tl]**
- Status and prospects for COHERENT

The First COHERENT Result: CsI[Na]

Led by U. Chicago group



J.I. Collar et al., NIM A773 (2016) 56-67

Scintillating crystal

- high light yield
- low intrinsic bg
- rugged and stable
- room temperature
- inexpensive

2 kg test crystal @U. Chicago.
Amcrys-H, Ukraine

Calibration of 14.6-kg detector at U. Chicago (^{241}Am, ^{133}Ba)

^{241}Am



^{133}Ba



Light yield:
13.35 pe/keVee,
uniform within ~2%

Used to determine
event selection efficiency

CsI quenching factor measurements at TUNL w/ neutrons



$$\underbrace{13.348\ \text{pe/keVee}}_{\text{ee light yield}} * \underbrace{0.0878\ \text{keVee/keVr}}_{\text{QF}} = \mathbf{1.2\ pe/keVr}$$

The CsI Detector in Shielding in Neutrino Alley at the SNS





A hand-held detector!

Almost wrapped up...

Layer	HDPE*	Low backg. lead	Lead	Muon veto	Water
Thickness	3”	2”	4”	2”	4”
Colour					

COHERENT data taking

1.76 x10^{23} POT delivered to CsI (7.48 GWhr)



Neutron background data-taking for ~2 years before first CEvNS detectors

CsI data-taking starting summer 2015

Example CsI waveform



- (C ROI) – (AC ROI) = CEvNS + Beam-on bg
- Pretraces used for afterglow background removal

Event Selection Cuts

Quality	Remove coincidences in muon veto, deadtime from PMT saturation blocking, digitizer range overflow	Select recoil-like low-energy pulses, reject muons
Afterglow	Reject signals with >=4 peaks (~spe) in pretrace	Remove afterglow (phosphorescence) contamination
"Cherenkov"	Require minimum number of peaks in the scintillation signal	Remove accidental coincidences between Cherenkov emission in PMT window and dark counts/ afterglow
Risetime	Pulse-shape based	Remove misidentified scintillator onset, accidental groupings of dark counts, etc.

- **2 independent analyses** with slightly different cut optimization yield consistent results
- "Analysis I" presented here

Event selection cut efficiencies



Data quality and stability: **fluctuations small and understood**



Energy to SNS target

CsI channel baseline

PMT SPE mean charge, used for gain fluctuation correction

Afterglow event removal fraction

Muon veto cut
Linear gate cut
DAQ overflow cut

Gain from internal crystal backgrounds

POT signal delay from muon panel neutron coincidences

Neutron backgrounds

- Evaluated using EJ-301 liquid scintillator cell deployed inside CsI shielding before CsI deployment
- Consistent with Geant4 simulation for SNS production & shielding





(consistent w/other measurements)

NINs: non-zero component at 2.9σ
(factor ~1.7 lower than prediction)

Expect: 0.93 ± 0.23 beam n events/GWhr

0.54 ± 0.18 NIN events/GWhr (neglected)

<~11 neutron events in CsI dataset

First light at the SNS with 14.6-kg CsI[Na] detector

Charge

Time



Observation of coherent elastic neutrino-nucleus scattering

D. Akimov[1,2], J. B. Albert[3], P. An[4], C. Awe[4,5], P. S. Barbeau[4,5], B. Becker[6], V. Belov[1,2], A. Brown[4,7], A. Bolozdy...
+ See all authors and affiliations

Science 03 Aug 2017:
eaao0990
DOI: 10.1126/science.aao0990

Peer Reviewed
← see details

D. Akimov et al., *Science*, 2017
http://science.sciencemag.org/content/early/2017/08/02/science.aao0990

Likelihood analysis: 2D in energy (pe) and time



$6 \leq PE \leq 30,\ 0 \leq t \leq 6000$ ns

Results of 2D energy, time fit

SM prediction, 173 events



Best fit: **134 ± 22** observed events

No CEvNS rejected at 6.7σ, consistent w/SM within 1σ

Signal, background, and uncertainty summary numbers

$6 \leq PE \leq 30$, $0 \leq t \leq 6000$ ns

Beam ON coincidence window	547 counts
Anticoincidence window	405 counts
Beam-on bg: prompt beam neutrons	7.0 ± 1.7
Beam-on bg: NINs (neglected)	4.0 ± 1.3
Signal counts, single-bin counting	136 ± 31
Signal counts, 2D likelihood fit	**134 ± 22**
Predicted SM signal counts	**173 ± 48**

Uncertainties on signal and background predictions	
Event selection	5%
Flux	10%
Quenching factor	25%
Form factor	5%
Total uncertainty on signal	**28%**
Beam-on neutron background	25%

Dominant uncertainty

What constraints do these data make on new interactions?

A first example: simple counting to constrain **non-standard interactions (NSI)** of neutrinos with quarks

Davidson et al., JHEP 0303:011 (2004)
Barranco et al., JHEP 0512:021 (2005)

"Model-independent" parameterization

$$\mathcal{L}_{\nu H}^{NSI} = -\frac{G_F}{\sqrt{2}} \sum_{\substack{q=u,d \\ \alpha,\beta=e,\mu,\tau}} [\bar{\nu}_\alpha \gamma^\mu (1-\gamma^5)\nu_\beta] \times (\varepsilon_{\alpha\beta}^{qL}[\bar{q}\gamma_\mu(1-\gamma^5)q] + \varepsilon_{\alpha\beta}^{qR}[\bar{q}\gamma_\mu(1+\gamma^5)q])$$

ε's parameterize new interactions

"Non-Universal": ε_{ee}, $\varepsilon_{\mu\mu}$, $\varepsilon_{\tau\tau}$

Flavor-changing: $\varepsilon_{\alpha\beta}$, where $\alpha \neq \beta$

$\Rightarrow$ some are quite poorly constrained (~unity allowed)

Cross-section for CEvNS including NSI terms

For flavor α, *spin zero* nucleus, and E<<k,M:

$$\left(\frac{d\sigma}{dE}\right)_{\nu N} = \frac{G_F^2 M}{\pi} F^2(2MT)\left[1 - \frac{MT}{2E_\nu^2}\right] \times$$

$$\{[Z(g_V^p + 2\varepsilon_{\alpha\alpha}^{uV} + \varepsilon_{\alpha\alpha}^{dV}) + N(g_V^n + \varepsilon_{\alpha\alpha}^{uV} + 2\varepsilon_{\alpha\alpha}^{dV})]^2$$ non-universal

$$+ \sum_{\alpha\neq\beta} [Z(2\varepsilon_{\alpha\beta}^{uV} + \varepsilon_{\alpha\beta}^{dV}) + N(\varepsilon_{\alpha\beta}^{uV} + 2\varepsilon_{\alpha\beta}^{dV})]^2\}$$ flavor-changing

$$\left.\begin{array}{r} g_V^p = (\frac{1}{2} - 2\sin^2\theta_W), \quad g_V^n = -\frac{1}{2} \\ \varepsilon_{\alpha\beta}^{qV} = \varepsilon_{\alpha\beta}^{qL} + \varepsilon_{\alpha\beta}^{qR} \end{array}\right\}$$ SM parameters

- NSI with these assumptions affect ***total cross-section*, *not differential shape of recoil spectrum***
- size of effect depends on N, Z
  (different for different elements)
- ε's can be negative and parameters can cancel

Ratio of rate with NSI to SM rate (all flavors in stopped-pion beam)

ε_{ee}^{uV} vs ε_{ee}^{dV} parameters (assume others zero)

CsI



Note that for

$$Z(g_V^p + 2\varepsilon_{ee}^{uV} + \varepsilon_{ee}^{dV}) + N(g_V^n + \varepsilon_{ee}^{uV} + 2\varepsilon_{ee}^{dV}) = \pm(Zg_V^p + Ng_V^n),$$

the rate is the same as for the SM, so parameters will be allowed

Get slightly different slope for different targets

Generalized mass ordering degeneracy in neutrino oscillation experiments

Pilar Coloma[1] and Thomas Schwetz[2]

Phys.Rev. D94 (2016) no.5, 055005,
Erratum: Phys.Rev. D95 (2017) no.7, 079903
Also: P. Coloma et al., JHEP 1704 (2017) 116



Normal ordering w/no NSI...



...looks just like inverted ordering w/NSI

If you allow for NSI to exist, you can't tell the neutrino mass ordering in long-baseline experiments

... NC scattering can constrain NSI...

➔DUNE may need this...

χ^2 fit results for current CsI data set: 90% allowed region



- Simple one-bin analysis
- Assume all other ε's zero

Separate bands not resolved due to current uncertainty (dominated by QF).. will improve, and different N targets will help

*CHARM constraints apply only to heavy mediators

A COHERENT enlightenment of the neutrino Dark Side

Pilar Coloma,[1,*] M. C. Gonzalez-Garcia,,[2,3,4,†] Michele Maltoni,,[5,‡] and Thomas Schwetz[6,§]



Global fits to COHERENT + oscillation experiments

Solid: COHERENT
Dashed: COHERENT + osc
Blue: LMA ($\theta_{12} < \pi/4$)
Red: LMA-D ($\theta_{12} > \pi/4$)
("dark side", still allowed with NSI)

1σ, 2σ allowed regions projected in (ε_{ee}^{uV}, $\varepsilon_{\mu\mu}^{uV}$) plane



Already meaningful constraints!

This is the first measurement of low-energy NC neutrino-hadron interaction with event-by-event *spectral information*



Low energy (<~100 MeV) NC measurements so far:

J.A. Formaggio and G. Zeller, RMP 84 (2012) 1307-1341

^{12}C excitation

15-MeV gamma observed

Isotope	Reaction Channel	Source	Experiment	Measurement (10^{-42} cm^2)	Theory (10^{-42} cm^2)
	$^{12}C(\nu_\mu,\nu_\mu)^{12}C^*$	Stopped π/μ	KARMEN	$3.2 \pm 0.5(\text{stat}) \pm 0.4(\text{sys})$	2.8 [CRPA] (Kolbe *et al.*, 1999b)
	$^{12}C(\nu,\nu)^{12}C^*$	Stopped π/μ	KARMEN	$10.5 \pm 1.0(\text{stat}) \pm 0.9(\text{sys})$	10.5 [CRPA] (Kolbe *et al.*, 1999b)

Deuterium breakup

$d(\bar{\nu}_e, \bar{\nu}_e)pn$ neutron counting

Experiment	Measurement	σ_{fission} (10^{-44} cm^2/fission)	$\sigma_{\text{exp}}/\sigma_{\text{theory}}$
Savannah River (Pasierb *et al.*, 1979)	$\bar{\nu}_e$NC	3.8 ± 0.9	0.8 ± 0.2
ROVNO (Vershinsky *et al.*, 1991)	$\bar{\nu}_e$NC	2.71 ± 0.47	0.92 ± 0.18
Krasnoyarsk (Kozlov *et al.*, 2000)	$\bar{\nu}_e$NC	3.09 ± 0.30	0.95 ± 0.33
Bugey (Riley *et al.*, 1999)	$\bar{\nu}_e$NC	3.15 ± 0.40	1.01 ± 0.13

That's it... (not many CC measurements in this range either)

Another phenomenological analysis, making use of spectral fit:

COHERENT constraints on nonstandard neutrino interactions

Jiajun Liao and Danny Marfatia

arXiv:1708.04255

SM weak charge

Effective weak charge in presence of light vector mediator Z'

$$Q^2_{\alpha,\mathrm{SM}} = (Zg_p^V + Ng_n^V)^2 \Rightarrow Q^2_{\alpha,\mathrm{NSI}} = \left[Z\left(g_p^V + \frac{3g^2}{2\sqrt{2}G_F(Q^2+M^2_{Z'})}\right) + N\left(g_n^V + \frac{3g^2}{2\sqrt{2}G_F(Q^2+M^2_{Z'})}\right)\right]^2$$

- Q^2-dependence ➔ *affects recoil spectrum*
- 2 parameters: g, $M_{Z'}$



Dashed: SM
Solid: NSI w/ $M_{Z'}$= 10 MeV, g=10^{-4}

Blue: ν_μ
Red: ν_μ + ν_μ–bar
Black: ν_μ + ν_μ–bar + ν_e



OUTLINE

- Coherent elastic neutrino-nucleus scattering (CEvNS)
- Why measure it? Physics motivations (short and long term)
- How to measure CEvNS
- The COHERENT experiment at the SNS
- **First light** with CsI[Tl]
- **Status and prospects for COHERENT**

What's Next for COHERENT?





Deployments so far in Neutrino Alley



COHERENT **CEvNS** Detector Status and Near Future

Nuclear Target	Technology	Mass (kg)	Distance from source (m)	Recoil threshold (keVr)	Data-taking start date
CsI[Na]	Scintillating Crystal	14.6	20	6.5	9/2015
Ge	HPGe PPC	10	22	5	2017
LAr	Single-phase	22	29	20	12/2016, upgraded summer 2017
NaI[Tl]	Scintillating crystal	185*/ 2000	28	13	*high-threshold deployment summer 2016





- CsI will continue running
- 185 kg of NaI installed in July 2016
  - taking data in high-threshold mode for CC on ^{127}I
  - PMT base modifications to enable low-threshold CEvNS running
- LAr single-phase detector installed in December 2016
  - upgraded w/TPB coating of PMT & Teflon, running since May 2017
- First Ge detectors to be installed late 2017

Single-Phase Liquid Argon

- ~22 kg fiducial mass
- 2 x Hamamatsu 5912-02-MOD 8” PMTs
  - 8” borosilicate glass windown
  - 14 dynodes
  - QE: 18%@ 400 nm
- Wavelength shifter: TB-coated teflon walls and PMTs
- Cryomech cryocooler – 90 Wt
  - PT90 single-state pulse-tube cold head



IU, UT, ORNL





Detector from FNAL, previously built (J. Yoo et al.) for CENNS@BNB
(S. Brice, Phys.Rev. D89 (2014) no.7, 072004)

Future LAr concepts

- 1-tonne scale feasible in Neutrino Alley
- Considering depleted argon
  to reduce ^{39}Ar background
- Considering SiPMs







High-Purity Germanium Detectors

P-type Point Contact



- Excellent low-energy resolution
- Well-measured quenching factor
- Reasonable timing

- Canberra cryostats in multi-port dewar
- Compact poly+Cu+Pb shield
- Muon veto
- Designed to enable additional detectors



- 10 kg of detectors available (MAJORANA unenriched prototypes)
- Under refurbishment/test at NCSU, Duke and LANL
- Dewar fabrication nearly complete
- Future: additional 2.5 kg detectors (UChicago, NCSU)

Sodium Iodide (NaI[Tl]) Detectors (NaIvE)

- up to 9 tons available, 2 tons in hand
- QF measured
- require PMT base refurbishment (dual gain) to enable low threshold for CEvNS on Na measurement
- development and instrumentation tests underway at UW, Duke

Multi-ton concept



In the meantime: **185 kg deployed at SNS** to go after ν_e**CC on** ^{127}**I**

Isotope	Reaction Channel	Source	Experiment	Measurement (10^{-42} cm^2)	Theory (10^{-42} cm^2)
^{127}I	^{127}I$(\nu_e, e^-)^{127}$Xe	Stopped π/μ	LSND	284 ± 91(stat) ± 25(sys)	210-310 [Quasi-particle] (Engel *et al.*, 1994)

J.A. Formaggio and G. Zeller, RMP 84 (2012) 1307-1341

COHERENT Non-CEvNS Detectors ("In-COHERENT")

Sandia Neutron Scatter Camera	Multiplane liquid scintillator	Neutron background	Deployed 2014-2016
SciBath	WLS fiber + liquid scintillator	Neutron background	Deployed 2015
NaI[Tl]	Scintillating crystal	ν_eCC	High-threshold deployment summer 2016
Lead Nube	Pb + liquid scintillator	NINs in lead	Deployed 2016
Iron Nube	Fe + liquid scintillator	NINs in iron	Deployed 2017
MARS	Plastic scintillator and Gd sandwich	Neutron background	Under deployment
Mini-HALO	Pb + NCDs	NINs in lead	In design






And many more ideas and activities for Neutrino Alley and beyond...

- Inelastic CC and NC in Ar, Pb, ...
- Other crystal or scint deployments in CsI shield
- Flux normalization using D_2O (well known xscn)
- Ancillary measurements: QF
- Directional detectors
- ...

Protons on target delivered so far





CEvNS

Neutron backgrounds

ν_eCC on ^{127}I

Neutron backgrounds

CEvNS

Neutrino-induced neutrons

Summary

- **First measurement of CEvNS** in COHERENT CsI[Na] in Neutrino Alley at the SNS
- Multiple physics motivations
  - DM bg, SM test, astrophysics, nuclear physics, ...
- Low-hanging fruit: **meaningful bounds on ν NSI**



- **It's just the beginning....**
- Multiple targets, upgrades and new ideas in the works!
- Other CEvNS experiments will soon join the fun
  (CONNIE, CONUS, MINER, RED, Ricochet, Nu-cleus...)

Extras/backups

Light DM direct detection possibilities

Light new physics in coherent neutrino-nucleus scattering experiments

Patrick deNiverville,[1] Maxim Pospelov,[1,2] and Adam Ritz[1]
[1] *Department of Physics and Astronomy, University of Victoria, Victoria, BC V8P 5C2, Canada*
[2] *Perimeter Institute for Theoretical Physics, Waterloo, ON N2J 2W9, Canada*
(Dated: May 2015)

production:

proton → target → $\pi^{0,\pm}$ →

$$\pi^0 \longrightarrow \gamma + V^{(*)} \longrightarrow \gamma + \chi^\dagger + \chi$$

$$\pi^- + p \longrightarrow n + V^{(*)} \longrightarrow n + \chi^\dagger + \chi$$

detection:







1 ton LAr
E_{rec}>20keVnr
10^{23} POT

R. Tayloe
Cosmic Visions 2017

NIN measurement in SNS basement with Nubes

Liquid scintillator surrounded by Pb, Fe (swappable for other NIN targets) inside water shield



P. Barbeau

Evaluation of 14.6-kg detector risetime-cut efficiency w/ ^{133}Ba data





Risetime cut applied to SNS data


Charge
Time
A
C (on)
AC (on)
B
C (on)
AC (on)
C
C (off)
AC (off)
D
C (off)
AC (off)
E
AC (on) - AC (off)
F
AC (on) - AC (off)
Counts / 2 PE
Counts / 500 ns
Number of photoelectrons (PE)
Arrival time (μs)

In-Situ bg limit on in-beam neutrons

Electron capture decay of ^{128}I at 31.8 keV

Inelastic scattering peak (57.6 keV) recoil + γ's

Neutron source outside shielding



90% CL maximum allowed neutron counts for Beam-ON data

Total residual counts vs time
consistent w/ entirely beam-induced events



χ^2 with pull for our situation, including background

(simple one-bin analysis)

$$\chi^2 = \frac{\left(N_{\rm meas} - N_{NSI}(\varepsilon_{ee}^{uV}, \varepsilon_{ee}^{dV})[1+\alpha] - B_{\rm on}[1+\beta]\right)^2}{\sigma_{\rm stat}^2} + \left(\frac{\alpha}{\sigma_\alpha}\right)^2 + \left(\frac{\beta}{\sigma_\beta}\right)^2 .$$

$N_{\rm meas}$ steady-state background-subtracted counts

$N_{\rm NSI}(\varepsilon_{ee}^{uV}, \varepsilon_{ee}^{dV})$ expected signal with NSI

$B_{\rm ss}$ expected steady-state background

$B_{\rm on}$ expected beam-on background

$$\sigma_{\rm stat} = \sqrt{N_{\rm meas} + 2B_{\rm SS} + B_{\rm on}}$$

$\sigma_{\rm sys,SS} = 0$ expected systematic on steady-state bg (assume zero because well measured)

α: for signal normalization systematic uncertainty

β: for beam-on background normalization uncertainty

SNS Beam Schedule

- 2100 hours @ 1 MW
- 1600 hours @ 1.2 MW



Production beam through September 30, 2017

SNS Beam Schedule

- 1100 hours @ 1.4 MW
- 5 month outage