

Unravelling the richness of dark sector by FASER ν

Yasaman Farzan

IPM, Tehran


This work is based on

- P. Bakhti, YF, S. Pascoli, “[Unravelling the richness of dark sector by FASER \$\nu\$](#) ,” arXiv: 2006.05437, to appear in JHEP

Work done during my stay at [ICTP](#), Trieste, Italy





LHC timeline and luminosity




Run III: 14 TeV; 150 fb^{-1} ; 2021-2023

Main detectors of the LHC




Smaller detectors

ZDC=Zero Degree Calorimeter




FASER

- FASER: ForwArd Search ExpeRiment




- FASER ν Run III: 14 TeV; 150 fb^{-1} ; 2021-2023



FASER collaboration, arXiv:1908.0231


10 m of concrete and 90 m of rock




FASER collaboration,
arXiv:1908.0231

$$Br(K^+ \rightarrow \mu^+ \nu_\mu) = 63\%$$

$$Br(K^+ \rightarrow \pi^0 e^+ \nu_e) = 5\%$$



FASER collaboration, arXiv:1908.0231



FASER collaboration, arXiv:1908.0231

25 cm × 25 cm × 1.35 m emulsion detector


tungsten target mass of 1.2 tons.

1300 ν_e , 20,000 ν_μ , and 20 ν_τ

New physics in FASER


- FASER Collaboration, [Phys. Rev. D 99 \(2019\) no.9, 095011](#);
- J. Alimena et al., [arXiv:1903.04497](#);
- J. L. Feng et al, [Phys. Rev. D 97 \(2018\) no.5, 055034](#);
- N. Okada and D. Raut, [arXiv:1910.09663](#);
- I. Boiarska et al, [arXiv:1908.04635](#);
- F. Kling and S. Trojanowski, [Phys. Rev. D 97 \(2018\) no.9, 095016](#);
- J. C. Helo, M. Hirsch and Z. S. Wang, [JHEP 1807 \(2018\) 056](#);
- F. Deppisch, S. Kulkarni and W. Liu, [Phys. Rev. D 100 \(2019\) 035005](#);
- J. L. Feng et al, [Phys. Rev. D 98 \(2018\) no.5, 055021](#);
- R. N. Mohapatra and N. Okada, [arXiv:1908.11325](#);
- A. Berlin and F. Kling, [Phys. Rev. D 99 \(2019\) no.1, 015021](#);
- K. Jodowski, F. Kling, L. Roszkowski and S. Trojanowski, [arXiv:1911.11346](#);
- D. Dercks, J. De Vries, H. K. Dreiner and Z. S. Wang, [Phys. Rev. D 99 \(2019\) no.5, 055039](#).

- FASER Collaboration, “FASER's Physics Reach for Long-Lived Particles,” Phys. Rev. D 99 (2019) no.9, 095011;



- FASER Collaboration, “FASER's Physics Reach for Long-Lived Particles,” Phys. Rev. D 99 (2019) no.9, 095011;

$$\frac{1}{2}m_X^2 X^\mu X_\mu - g_X j_\mu^X X^\mu - \frac{\epsilon}{2 \cos \theta_W} B_{\mu\nu} X^{\mu\nu}$$










Testing 3+1 model at FASER ν

$$P(\nu_\alpha \rightarrow \nu_\alpha) = 1 - 4|U_{\alpha 4}|^2(1 - |U_{\alpha 4}|^2) \sin^2 \frac{\Delta m_{41}^2 L}{4E}$$
$$P(\nu_\alpha \rightarrow \nu_\beta) = \sin^2 2\theta_{\alpha\beta} \sin^2 \frac{\Delta m_{41}^2 L}{4E}.$$

FASER collaboration, arXiv:1908.0231

$$m_{\nu_4} = 40 \text{ eV}$$






New physics with FASER ν

- M. Bahraminasr, P Bakhti and M Rajae, “Sensitivities to secret neutrino interaction at FASER ν ,” arXiv:2003.09985;
- Felix Kling, “Probing Light Gauge Bosons in Tau Neutrino Experiments,” arXiv: 2005.03594.
- P. Bakhti, YF, S. Pascoli, “Unravelling the richness of dark sector by FASER ν ,” arXiv: 2006.05437, to appear in JHEP

Parton Distribution Function (PDF)



Two partons colliding

$$p_1 = p(x_1, 0, 0, x_1) \quad \text{and} \quad p_2 = p(x_2, 0, 0, -x_2)$$

$$s = 4x_1 x_2 p^2$$

$$p = 7 \text{ TeV}$$

$$x_1 \sim 0.1, \ x_2 \sim 10^{-7} \quad \longrightarrow \quad s \sim \text{GeV}^2$$

Kinematics tells us that particles lighter than GeV can be produced and emitted in the **forward** direction.

Need for intermediate particles

- At the **Interaction Point (IP)**



- X' should not have too small coupling to SM particles to be abundantly produced at IP.



- X should be feebly interacting so that survive up to FASER ν .

Signature of X particles at FASER ν .

$$X \rightarrow f + \bar{f} \longrightarrow \text{Standard model fermions}$$

$$X \rightarrow f + \bar{f} + Y \longrightarrow \text{Missing energy-momentum; dark matter?}$$

Decay length 1 mm-10 cm

$$X \rightarrow \eta + \bar{\eta} \quad \eta \rightarrow f\bar{f} \quad \bar{\eta} \rightarrow f\bar{f}$$

$$X \rightarrow \eta + \bar{\eta} + Y \quad \eta \rightarrow f\bar{f} \quad \bar{\eta} \rightarrow f\bar{f}$$


Dark matter

What is dark matter made of?

WIMP paradigm in its heyday

- WIMP=Weakly Interacting Massive Particle

Freeze-out scenario of dark matter



$$\langle v_{DM} \sigma(DM + DM) \rangle = 1 \text{ pb}$$

$$\langle v_{DM} \sigma(DM + DM) \rangle \sim \frac{g^4}{4\pi M^2}$$

$$g \sim e \sin \theta_W$$



$$M \sim \text{few} \times 100 \text{ GeV}$$


Hopes for WIMP discovery

- LHC: $p+p \rightarrow DM + \text{monophoton}$ or $DM + \text{monojet}$
- Direct DM searches: $DM + \text{nucleus} \rightarrow DM + \text{recoiled nucleus}$
- Indirect DM searches: $DM + DM \rightarrow SM \text{ particles}$

In regions that DM density is higher than the average: Galaxy center, DM halo, Earth or sun center

High hopes for DM detection

Null results so far




<http://newscenter.lbl.gov>


In 2013, WIMP was still **young**.

Italian: WIMP era ancora **giovane**

Iranian: **Javan**



E. Aprile et al. (XENON1T), Phys. Rev. Lett. 121 (2018) 111302.




<http://newscenter.lbl.gov>

In 2013, WIMP was still **young**.

Italian: WIMP era ancora **giovane**

Iranian: **Javan**

Iranian saying



آرزو بسر جوانان بود عیب، ولی
عیب ایجاد است که امید جوان رفته به باد!

محمد شیران (عاصم اصفهانی)

High hopes for DM detection



*Hope for WIMP is **not** given up.*

*But the Hegemony of WIMP as **THEE DM candidate** is broken*

Time to leave the comfort zone


- Comfort zone: 100 GeV-1 TeV
- Whole range: $M > 10^{-21} eV$

GeV scale DM



CDEX collaboration, arXiv: 2007.15555.

More precise evaluation of annihilation from DM abundance



Freeze-out scenario

$$\sigma \sim 1 \text{ pb}$$

Steigman, Dasgupta and Beacom, *Phys.Rev.D* 86 (2012) 023506

Light dark matter annihilation rate

$$\sigma \sim \frac{g^4}{4\pi M^4} m_{DM}^2$$



M= Mass of the mediator

$$m_{DM} \sim \text{GeV}$$

If annihilation is to SM particles: electron, muon, u, d, s, gluons or photons

$$M/g > \text{TeV} \quad \longrightarrow \quad \sigma \ll pb$$

One exception: SLIM model


- New fields:
- Majorana Right-handed neutrino
- SLIM=Scalar as Light as MeV

Boehm, Y. F., T. Hambye, S. Palomares-Ruiz and S. Pascoli, PRD 77 (2008) 43516;

- Effective Lagrangian: $\mathcal{L}_I \supset g\phi\bar{N}\nu$

- New parameters:
 $g \quad m_\phi \quad m_N$

Annihilation cross-section




$$\begin{aligned}\langle \sigma(\phi\phi \rightarrow \nu\nu)v_r \rangle &= \langle \sigma(\phi\phi \rightarrow \bar{\nu}\bar{\nu})v_r \rangle \\ &\simeq \frac{g^4}{4\pi} \frac{m_N^2}{(m_\phi^2 + m_N^2)^2},\end{aligned}$$

$$g \simeq 10^{-3} \sqrt{\frac{m_N}{10 \text{ MeV}}} \left(\frac{\langle \sigma v_r \rangle}{10^{-26} \text{ cm}^3/\text{s}} \right)^{1/4} \left(1 + \frac{m_\phi^2}{m_N^2} \right)^{1/2}$$

neutrino masses

- In this scenario, SLIM does not develop any VEV so the tree level neutrino mass is zero.
- Radiative mass in case of real SLIM scalar:



Ultraviolet cutoff Λ

$$m_\nu = \frac{g^2}{16\pi^2} m_N \left[\ln\left(\frac{\Lambda^2}{m_N^2}\right) - \frac{m_\phi^2}{m_N^2 - m_\phi^2} \ln\left(\frac{m_N^2}{m_\phi^2}\right) \right]$$

Bounds on SLIM mass

$$m_\phi < M_N$$

$$O(1) \text{ MeV} \lesssim m_N \lesssim 10 \text{ MeV}.$$

GeV scale Dark matter

$$DM + DM \rightarrow \eta + \bar{\eta}$$

Beyond SM

$$\sigma \sim \frac{g^4}{4\pi M^4} m_{DM}^2$$

$$m_{DM} \sim \text{GeV}$$

$$M/g \sim 100 \text{ GeV}$$



$$\sigma \sim 1 \text{ pb}$$

GeV scale Dark matter

$$DM + DM \rightarrow \eta + \bar{\eta}$$

Beyond SM

$$m_{DM} \sim \text{GeV}$$


$$m_\eta < m_{DM}$$

$$\eta \rightarrow f\bar{f}$$


SM fermion

$$\left. \begin{array}{l} m_\eta > 10 \text{ MeV} \\ \tau_\eta < 1 \text{ sec} \end{array} \right\} \xrightarrow{\quad} \text{No trouble for BBN}$$

Shift of paradigm in DM model building



Inspired by 5% of matter (ordinary matter)



Richness of SM makes life dazzling




- Why should DM be minimal?
- It would be boring if it is so.
- More exciting possibilities: Multicomponent DM, SIDM,....

Can we uncover the richness of dark sector?

- We should **not** overlook any possibility to do so.
- FASER ν : Superb **position** and **spatial** resolution


Reconstructing **invisible** tracks of **neutral** particles decaying to charged leptons
- The point of our paper: FASER ν can be a useful tool.



FASER collaboration, arXiv:1908.0231

10 m of concrete and 90 m of rock

Toy model

- Intermediate X' is produced at IP.

- Before reaching detector, $X' \rightarrow X\bar{X}$


- X decays at FASER ν to the SM charged leptons:

$$X \rightarrow \eta\bar{\eta}Y \rightarrow \text{Missing energy and momentum}$$

- η and $\bar{\eta}$ decay after 1mm-10cm: $\eta \rightarrow l\bar{l}$ $\bar{\eta} \rightarrow l\bar{l}$


A generic scenario

Specific model
Demonstrating ability
Of FASER ν



Signature at FASER ν


- Two pairs charged leptons plus missing energy



Energy momentum of leptons can be measured



The energy momentum of $\bar{\eta}$ and η



Need for intermediate particles

- At the **Interaction Point (IP)**



- X' should not have too small coupling to the SM particles to be abundantly produced at IP.



Possibilities for X'

- 1) Z' boson of $B - a_e L_e - a_\mu L_\mu - a_\tau L_\tau$ (with $a_e + a_\mu + a_\tau = -3$) or kinetically mixed $\epsilon Z'_{\mu\nu} B^{\mu\nu}$
- 2) Scalar ϕ coupled to quarks: $\phi \bar{u}u, \phi \bar{d}d, \phi \bar{s}s$

Mixed with SM Higgs or Mixed with new inert Higgs

Small mixing to first generation

Production from Heavy meson decay at IP


arbitrary coupling to first generation

Direct production from parton interaction at IP

Possibilities for X'

- 1) Z' boson of $B - a_e L_e - a_\mu L_\mu - a_\tau L_\tau$ (with $a_e + a_\mu + a_\tau = -3$) or kinetically mixed $\epsilon Z'_{\mu\nu} B^{\mu\nu}$
- 2) Scalar ϕ coupled to quarks: $\phi \bar{u}u, \phi \bar{d}d, \phi \bar{s}s$
- 3) Pseudoscalar axion:
$$\frac{X' G_{\mu\nu}^i G_{\alpha\beta}^i \epsilon^{\mu\nu\alpha\beta}}{\Lambda},$$

solving the QCD θ -term.



$$\Lambda = 8\pi f_a / \alpha_{QCD}$$

Fast decay of X'

- X' with a mass larger than 3 pion mass decays fast to hadrons
- We want it to decay faster to $X \bar{X}$

Scalar X : $\lambda m_{X'} X' \bar{X} X$

Fermion X : $\lambda X' \bar{X} X.$

$$\lambda > 10^{-6}$$



$$Br(X' \rightarrow X \bar{X}) \gg Br(X' \rightarrow \text{hadrons})$$

X' will decay after traveling less than one meter.

Production of X'

Interaction Point (IP): $g(p_1) + g(p_2) \rightarrow X'(p_1 + p_2)$

$$p_1 = p(x_1, 0, 0, x_1) \quad \text{and} \quad p_2 = p(x_2, 0, 0, -x_2)$$

$$m_{X'} = p\sqrt{4x_1x_2} \quad \text{where} \quad \frac{m_{X'}^2}{4p^2} < x_1, x_2 < 1.$$

$$\langle |M^2| \rangle = \frac{2p^4 x_1^2 x_2^2}{\Lambda^2}.$$

$$\frac{d\sigma_{tot}(E_{X'})}{dE_{X'}} = \frac{\pi}{32\Lambda^2} \frac{m_{X'}^2}{p^2} \frac{1}{P_{X'}} F_g\left(\frac{E_{X'} + P_{X'}}{2p}\right) F_g\left(\frac{E_{X'} - P_{X'}}{2p}\right)$$

Production of X'

Interaction Point (IP): $g(p_1) + g(p_2) \rightarrow X'(p_1 + p_2)$

$$p_1 = p(x_1, 0, 0, x_1) \quad \text{and} \quad p_2 = p(x_2, 0, 0, -x_2)$$

$$m_{X'} = p\sqrt{4x_1x_2} \quad \text{where} \quad \frac{m_{X'}^2}{4p^2} < x_1, x_2 < 1.$$

$$\frac{d\sigma_{tot}(E_{X'})}{dE_{X'}} = \frac{\pi}{32\Lambda^2} \frac{m_{X'}^2}{p^2} \frac{1}{P_{X'}} F_g\left(\frac{E_{X'} + P_{X'}}{2p}\right) F_g\left(\frac{E_{X'} - P_{X'}}{2p}\right)$$

Gluon PDFs at $Q^2 = m_{X'}^2$

Angular distribution of the emitted X'

Interaction Point (IP): $g(p_1) + g(p_2) \rightarrow X'(p_1 + p_2)$

Transverse momentum ignored

$$p_t = 200 \text{ MeV}$$

$$p_1 = p(x_1, 0, 0, x_1) \quad \text{and} \quad p_2 = p(x_2, 0, 0, -x_2)$$

emission of X' within a small solid angle of $\pi\theta_t^2$ where $\theta_t = p_t/E_{X'}$

Transverse momentum of protons in a bunch $< 1 \text{ MeV} \ll 200 \text{ MeV}$

Is this the only production mode?

$$g + g \rightarrow X' + g \text{ or } g + q \rightarrow X' + q.$$

High transverse momentum for X'



Reaching ATLAS (and CMS) but not FASER


X particles directed towards FASER:

$$dN(E_X)/dE_X|_{2 \rightarrow 2} < 20\% \text{ of } dN(E_X)/dE_X|_{2 \rightarrow 1}$$

Angular spread of X

Momentum of X in the rest frame of X' : $k = (m_{X'}^2/4 - m_X^2)^{1/2}$


X' velocity in the lab frame: $v_{X'} = (1 - m_{X'}^2/E_{X'}^2)^{1/2}$




$$\theta_s = \arctan \left(\frac{2k}{E_{X'} v_{X'} \sqrt{1 - 4k^2/(m_{X'}^2 v_{X'}^2)}} \right)$$

For $p_t \ll k \sim m_{X'}/2$,


For $m_X \rightarrow m_{X'}/2$, $k \ll m_X/2$



fraction of X reaching FASER ν




$$\theta_d = 5 \times 10^{-4}$$



$$\theta_s = \arctan \left(\frac{2k}{E_{X'} v_{X'} \sqrt{1 - 4k^2 / (m_{X'}^2 v_{X'}^2)}} \right)$$

For $p_t \ll k \sim m_{X'}/2$,

For $m_X \rightarrow m_{X'}/2$, $k \ll m_X/2$




$$f = \theta_d^2 / (2(1 - \cos \theta_s))$$

$$f = \theta_d^2 / \theta_t^2$$

Spectrum of X from X' decay at FASER

Momentum of X in the rest frame of X' : $k = (m_{X'}^2/4 - m_X^2)^{1/2}$


X' velocity in the lab frame: $v_{X'} = (1 - m_{X'}^2/E_{X'}^2)^{1/2}$



$$\theta_s = \arctan \left(\frac{2k}{E_{X'} v_{X'} \sqrt{1 - 4k^2/(m_{X'}^2 v_{X'}^2)}} \right)$$

For $p_t \ll k \sim m_{X'}/2$,

For $m_X \rightarrow m_{X'}/2$, $k \ll m_X/2$



$$E_X = \frac{E_{X'}}{2} \left(1 + \frac{2k}{m_{X'}} \right)$$

Spectrum of X from X' decay at FASER and FASER ν

$$\frac{d\sigma_{tot}(E_{X'})}{dE_{X'}} = \frac{\pi}{32\Lambda^2} \frac{m_{X'}^2}{p^2} \frac{1}{P_{X'}} F_g\left(\frac{E_{X'} + P_{X'}}{2p}\right) F_g\left(\frac{E_{X'} - P_{X'}}{2p}\right)$$

$$\frac{dN(E_X)}{dE_X} = L f \frac{d\sigma_{tot}}{dE_{X'}}|_{E_X} \frac{2}{1 + \sqrt{1 - 4m_X^2/m_{X'}^2}}$$

Luminosity

$$E_X = \frac{E_{X'}}{2} \left(1 + \frac{2k}{m_{X'}}\right) \quad dE_{X'}/dE_X$$

Spectrum of X emitted towards FASER and FASER ν


$$\frac{d\sigma_{tot}(E_{X'})}{dE_{X'}} = \frac{\pi}{32\Lambda^2} \frac{m_{X'}^2}{p^2} \frac{1}{P_{X'}} F_g\left(\frac{E_{X'} + P_{X'}}{2p}\right) F_g\left(\frac{E_{X'} - P_{X'}}{2p}\right)$$

$$\frac{dN(E_X)}{dE_X} = Lf \frac{d\sigma_{tot}}{dE_{X'}}|_{E_X} \frac{2}{1 + \sqrt{1 - 4m_X^2/m_{X'}^2}}$$



$$f = \theta_d^2 / (2(1 - \cos \theta_s)) \propto \text{detector area}$$

Spectrum of X emitted towards FASER ν



run III of LHC


$\sqrt{s} = 14 \text{ TeV}$

150 fb^{-1}

$m_{X'} = 3 \text{ GeV}, m_X = 1 \text{ GeV}$


$\Lambda = 2 \times 10^4 \text{ GeV}$

Spectrum of X emitted towards FASER ν



PDFs: J. Rojo *et al.*, J. Phys. G **42** (2015) 103103

Spectrum of X emitted towards FASER ν



Total number = O(100000)

run III of LHC

$\sqrt{s} = 14 \text{ TeV}$

150 fb^{-1}

$m_{X'} = 3 \text{ GeV}, m_X = 1 \text{ GeV}$

$\Lambda = 2 \times 10^4 \text{ GeV}$

Number of X decaying at FASER ν

$$\frac{dN(E_X)}{dE_X} \exp^{-\frac{d}{\tau_X \gamma_X}} \left(\frac{s_z}{\tau_X \gamma_X} \right) dE_X$$

\propto detector area

Detector length

Proportional to Volume

Number of X decaying at FASER ν

$$\frac{dN(E_X)}{dE_X} \exp^{-\frac{d}{\tau_X \gamma_X}} \left(\frac{s_z}{\tau_X \gamma_X} \right) dE_X$$

$$\gamma_X = m_X/E_X$$


Length of the detector:

$\left[\begin{array}{l} \text{FASER}\nu \\ \text{FASER2: (upgrade of FASER for HL - LHC)} \end{array} \right]$


$$s_z = 1.3 \text{ m}$$

$$s_z = 5 \text{ m}$$

Bound on the coupling



Bound on the coupling




$$\frac{dN(E_X)}{dE_X} \exp^{-\frac{d}{\tau_X \gamma_X}} \left(\frac{s_z}{\tau_X \gamma_X} \right) dE_X$$

$$\tau_X \Lambda^2 = cte$$

$m_{X'} = 3 \text{ GeV}$ and $m_X = 1 \text{ GeV}$.

Spectrum of X emitted towards FASER ν



Total number = O(100000)

8/27/20

$$\frac{dN(E_X)}{dE_X} \exp^{-\frac{d}{\tau_X \gamma_X}} \left(\frac{s_z}{\tau_X \gamma_X} \right) dE_X$$

<0.01

run III of LHC

$\sqrt{s} = 14 \text{ TeV}$

150 fb^{-1}

$m_{X'} = 3 \text{ GeV}, m_X = 1 \text{ GeV}$

$\Lambda = 2 \times 10^4 \text{ GeV}$

Number of events < or = 100

71

Toy model

- Intermediate X' is produced at IP.

- Before reaching detector, $X' \rightarrow X\bar{X}$

- X decays at FASER ν to the SM charged leptons:

$$X \rightarrow \eta\bar{\eta}Y$$

Missing energy and momentum


- η and $\bar{\eta}$ decay after 1mm-10cm: $\eta \rightarrow l\bar{l}$ $\bar{\eta} \rightarrow l\bar{l}$

A generic scenario
NO BACKGROUND

Specific model
Demonstrating ability
Of FASER ν

Signature at FASER ν

- Two pairs charged leptons plus missing energy



$$\theta_\eta = m_X/E_X$$

$$\theta_l \sim \sqrt{m_\eta^2 - 4m_l^2}/E_\eta$$

$$\Delta\theta_\eta \sim \sqrt{(\Delta E/E)^2\theta_l^2/2 + 2\sigma_{ang}^2},$$

$$\sigma_{pos} = 0.4 \text{ } \mu\text{m}$$

$$\sigma_{ang} = \sqrt{2}\sigma_{pos}/L_{tr}$$

$$L_{tr} = 1 \text{ cm}$$

$$\sigma_{angle} = 0.06 \text{ mrad.}$$

$$\Delta E/E \sim 30\%.$$

$$X \rightarrow \eta + \bar{\eta} + Y \text{ or } X \rightarrow \eta + \bar{\eta}$$

$$\Delta P_t < |\vec{p}_\eta^t + \vec{p}_{\bar{\eta}}^t|$$


$$\Delta P_t = \left(2(\frac{\Delta E}{E})^2 [(p_\eta^t)^2 + (p_{\bar{\eta}}^t)^2] + (\sum |\vec{p}_l|^2 + \sum |\vec{p}_{\bar{l}}|^2) \sigma_{ang}^2 + |\vec{p}_\eta + \vec{p}_{\bar{\eta}}|^2 (\Delta \hat{z})^2 \right)^{1/2}$$

$$(\Delta E/E)^2 \sim 0.3^2 \quad (\sum |\vec{p}_l|^2 + \sum |\vec{p}_{\bar{l}}|^2) \sim |\vec{p}_\eta|^2 \sim |\vec{p}_{\bar{\eta}}|^2 \quad \Delta \hat{z} \sim 10^{-3}$$

$$X \rightarrow \eta + \bar{\eta} + Y \text{ or } X \rightarrow \eta + \bar{\eta}$$

$$\Delta P_t < |\vec{p}_\eta^t + \vec{p}_{\bar{\eta}}^t|$$

$$\Delta P_t = \left(2\left(\frac{\Delta E}{E}\right)^2 [(p_\eta^t)^2 + (p_{\bar{\eta}}^t)^2] + (\sum |\vec{p}_l|^2 + \sum |\vec{p}_{\bar{l}}|^2) \sigma_{ang}^2 + |\vec{p}_\eta + \vec{p}_{\bar{\eta}}|^2 (\Delta \hat{z})^2 \right)^{1/2}$$



Angle between projections of momenta to **xy** plane

$< 90^\circ$

+

$> 90^\circ$

-


Reconstructing the mass of η

$$m_\eta^2 = (p_l + p_{\bar{l}})^2$$

$$\Delta m_{sin}^2 / m_\eta^2 \sim [2(\Delta E/E)^2 + 2(\sigma_{ang}/\theta_l)^2]^{1/2} = 40\%$$

N pairs: $\Delta m_{sin}^2 / \sqrt{N}$  4 %

Decay length of η



$$\delta L_z \sim [(l_\eta^2 + l_{\bar{\eta}}^2)(\Delta\theta_\eta/\theta_\eta)^2 + 2\sigma_{pos}^2/\theta_\eta^2]^{1/2}$$


$1.5 \text{ cm} < l_\eta, l_{\bar{\eta}} < \text{few } 10 \text{ cm},$

$$\bar{\tau}_\eta = \frac{\sum_{i=1}^N l_\eta^i m_\eta / E_\eta}{N}$$

$$\frac{\delta\tau^{stat}}{\bar{\tau}} = \left(\frac{2 \log 2}{N}\right)^{1/2} \xrightarrow{\hspace{1cm}} 10 \%$$

Determining the masses of X and Y

- We do not know the energy of initial X



Any information on the mass of X and Y is limited by both **statistics** and **uncertainties** on **PDF**

Unlike m_η

Connection to dark matter

- X' can be produced in the early universe by gluon interaction.

For $T > m_{X'}$ and $\Lambda \lesssim 10^8 \text{ GeV}$

- X' decay can produce X and then Y (DM).

Connection to DM

Fermionic X and Y:

$$g_Y \eta \bar{V} Y + g_X \eta \bar{V} X$$

Complex

$$\sigma(Y + \bar{Y} \rightarrow \eta + \bar{\eta}) = \frac{g_Y^4}{m_V^4} \frac{m_Y^2}{4\pi}$$

$$\sigma(Y + \bar{Y} \rightarrow \eta + \bar{\eta}) \sim \text{pb}$$



$$\frac{m_V}{g_Y} \sim 10 \text{ GeV} \left(\frac{m_Y}{0.1 \text{ GeV}} \right)^{1/2}$$

For real case, cancelation between u and t channels

$$m_V > m_X \longrightarrow X \cancel{\rightarrow} V \bar{\eta}$$

$$\Gamma(X \rightarrow Y \eta \bar{\eta}) \sim \frac{g_Y^2 g_X^2}{100\pi^3} \frac{m_X^5}{m_V^4}$$

8/27/20 $\Gamma(X \rightarrow Y \eta \bar{\eta})(m_X/E_X) \sim (480 \text{ m})^{-1}$



$$g_X \sim 10^{-5} \left(\frac{\text{GeV}}{m_X} \right)^2 \left(\frac{m_V}{m_X} \right) \left(\frac{E_X}{500 \text{ GeV}_{80}} \right)$$

Connection to DM

Fermionic X and Y:

$$g_Y \eta \bar{V} Y + g_X \eta \bar{V} X$$

Complex

$$\sigma(Y + \bar{Y} \rightarrow \eta + \bar{\eta}) = \frac{g_Y^4}{m_V^4} \frac{m_Y^2}{4\pi}$$


$$\sigma(Y + \bar{Y} \rightarrow \eta + \bar{\eta}) \sim \text{pb}$$



$$\frac{m_V}{g_Y} \sim 10 \text{ GeV} \left(\frac{m_Y}{0.1 \text{ GeV}} \right)^{1/2}$$

For real case, cancelation between u and t channels

$$m_V > m_X$$




$$Z_2 \times Z_2$$

Exact

η and V are odd

X and Y odd




Connection to DM

Scalar X and Y:

$$\lambda_1 \bar{Y} X \bar{\eta} \eta + \lambda_2 \bar{X} X \bar{\eta} \eta + \lambda_3 \bar{Y} Y \bar{\eta} \eta.$$

Real or complex



Connection to DM

Scalar X and Y:

$$\lambda_1 \bar{Y} X \bar{\eta} \eta + \lambda_2 \bar{X} X \bar{\eta} \eta + \lambda_3 \bar{Y} Y \bar{\eta} \eta.$$

$$\Gamma(X \rightarrow Y \bar{\eta} \eta) \sim \frac{\lambda_1^2 m_X}{100 \pi^3}$$

$$\Gamma(X \rightarrow Y \bar{\eta} \eta)(m_X/E_X) \sim 1/(480 m),$$

$$\lambda_1 = 8 \times 10^{-7} (\text{GeV}/m_X) (E_X/500 \text{ GeV})^{1/2}$$

Connection to DM

Scalar X and Y:

$$\lambda_1 \bar{Y} X \bar{\eta} \eta + \lambda_2 \bar{X} X \bar{\eta} \eta + \lambda_3 \bar{Y} Y \bar{\eta} \eta.$$

$$\lambda_1 = 8 \times 10^{-7} (\text{GeV}/m_X) (E_X/500 \text{ GeV})^{1/2}$$

$$\sigma(Y\bar{Y} \rightarrow \eta\bar{\eta}) \sim \lambda_3^2 / (4\pi m_Y^2) \simeq 1 \text{ pb},$$



$$\lambda_3 \sim 10^{-4} \sqrt{m_Y/\text{GeV}}$$

$$\lambda_1 \ll \lambda_3$$

Conclusion

- FASER ν , with its superb abilities to reconstruct tracks, is ideal to study new long-lived GeV-scale feebly interacting particles that go through chain decays.
- Intriguing possibility for exploring GeV scale dark matter

Eta decay

$$\lambda_\eta \eta \bar{l}l. \quad \longrightarrow \quad \eta \rightarrow l\bar{l}.$$

$$\tau_\eta = 4\pi/(m_\eta \lambda_\eta^2)$$

$$\lambda_\eta > 10^{-7}, \quad \tau_\eta E_\eta / m_\eta < 1 \text{ m}$$

$$\eta H^\dagger \Phi + \eta^\dagger \Phi^\dagger H$$

